



Other Formats: [Citation](#) [MEDLINE](#)

Links: [Related Articles](#)

☐ Order this document

Pancreas 1998 Nov;17(4):325-33

Current concepts in pancreatic cancer: symposium summary.

Dugan MC, Sarkar FH

Pathologists Medical Group, Little Company of Mary Hospital, Torrance, California, USA.

This recent symposium featured speakers from several clinical and research disciplines. Among the findings: peptic ulcer disease is a significant predisposing risk factor (odds ratio = 3.9) for pancreatic cancer; as many as 50% of all intraductal papillary mucinous neoplasms are associated with invasive adenocarcinomas; alteration of gene expression via methylation of a gene promotor region constitutes a potentially reversible method of tumor suppressor gene inactivation; > 400 transcriptional alterations of gene expression have been identified for pancreatic cancer; some common molecular markers such as p53 and HER-2/neu may be related to morphologic alterations of in situ neoplasia and to transcriptional alterations of gene expression rather than mutational events; epidermal growth factor (EGF), transforming growth factor beta (TGF-beta), and related molecules may modulate gene transcription via "autocrine" or "paracrine" mechanisms; several cytokines, amylin (islet amyloid polypeptide), and other cachexia factors are responsible for paraneoplastic peripheral insulin resistance, ineffective utilization of glucose, and profound cachexia. In the clinical diagnostic arena: the World Health Organization established a standard nomenclature for intraductal papillary mucinous neoplasms, mucinous cystic tumors, intraductal mucinous hyperplasias, and solid pseudopapillary tumors; focal glandular differentiation may be commonly identified within pancreatic endocrine neoplasms (islet cell tumors) while not necessarily implying an unfavorable prognosis typical of ductal adenocarcinomas; positron emission tomography scanning may be used for evaluation of early tumor response to novel chemotherapeutic regimens; helical computed tomography (CT) is the state of the art in preoperative imaging for pancreatic cancer; neoadjuvant 5-fluorouracil (5-FU)-based chemoradiation in 39 "resectable" patients provided a median survival of 19 months, actuarial 4-year survival of 19%, and improved local tumor control; gemcitabine has shown promise in alleviating tumor-related symptoms with a significantly better "clinical benefit response" than single agent 5-FU (23.8 vs. 4.8%, $p = 0.0022$) based on change in pain intensity, daily analgesic consumption, performance status, and weight; a significant survival advantage was demonstrated in patients treated with conventional therapies whose tumors expressed p21WAF-1, an important inhibitor of cell cycle progression and downstream molecule of p53 and TGF-beta; a p21-adenovirus (rAD-p21) gene therapy resulted in significant growth inhibition of pancreatic cancer cell lines in tissue culture, and development of a successful SCID mouse-human pancreatic adenocarcinoma xenograft model provided an animal model for preclinical trials of rAD-p21.

Publication Types: